

This copy is for your personal, non-commercial use only.

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

Permission to republish or repurpose articles or portions of articles can be obtained by following the guidelines [here](#).

The following resources related to this article are available online at www.sciencemag.org (this information is current as of November 6, 2014):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/content/346/6210/1257521.full.html>

Supporting Online Material can be found at:

<http://www.sciencemag.org/content/suppl/2014/11/05/346.6210.1257521.DC1.html>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/content/346/6210/1257521.full.html#related>

This article **cites 53 articles**, 21 of which can be accessed free:

<http://www.sciencemag.org/content/346/6210/1257521.full.html#ref-list-1>

This article has been **cited by** 2 articles hosted by HighWire Press; see:

<http://www.sciencemag.org/content/346/6210/1257521.full.html#related-urls>

This article appears in the following **subject collections**:

Cell Biology

http://www.sciencemag.org/cgi/collection/cell_biol

RESEARCH ARTICLE SUMMARY

LOCAL TRANSLATION

Principles of ER cotranslational translocation revealed by proximity-specific ribosome profiling

Calvin H. Jan, Christopher C. Williams, Jonathan S. Weissman*

INTRODUCTION: Localized protein synthesis plays a critical role in creating subcellular structures by allowing protein production at the site of action and in response to local cellular need. Local translation is involved in diverse processes, including developmental patterning, cellular motility, synaptic plasticity, and protein trafficking through the secretory pathway. Despite this broad importance, few gene expression tools are available that faithfully preserve spatial information. We developed a flexible deep sequencing-based methodology (termed proximity-specific ribosome profiling) that enables precise characterization of localized protein synthesis. We applied our method to analyze translation at the endoplasmic reticulum (ER) in yeast and mammalian cells.

RATIONALE: The basis of our approach is to biotinylate ribosomes in intact cells in a manner dependent on their subcellular location. This is accomplished through the coexpression of a spatially restricted biotin ligase (BirA) fusion protein together with ribosomes containing an AviTag, which makes them substrates for BirA. Controlled pulses of biotin are then provided to allow for spatiotemporal control of ribosome labeling. This *in vivo* biotinylation enables the recovery of ribosomes from defined locations, including those that cannot be purified by classical cell fractionation techniques. Combining this purification strategy with ribosome profiling, the deep sequencing of ribosome-protected mRNA fragments, provides subcodon resolution of which messages were translated at the site of interest.

RESULTS: We identified several principles used by cells to coordinate translation with ER targeting.

Cotranslational targeting to the ER is pervasive and is principally determined by the location of the hydrophobic targeting sequence within the protein, rather than the mechanism of targeting or translocation. The position of this hydrophobic domain within the open reading frame determines the duration of time a targeted ribosome nascent-chain complex (RNC) can associate with the ER. Our data suggest a role for polysomes in retaining mRNAs at the ER, allowing for efficient targeting of RNCs for translocation.

Position-specific analyses revealed that distinct translocon complexes engage nascent chains at different points during synthesis. Most proteins engage the ER immediately after or even before the signal sequence or signal anchor emerges from the ribosome. These nascent chains typically undergo a conformational rearrangement within the translocon, the proteinaceous tunnel through which nascent proteins cross the ER membrane. This rearrangement results in a “looped” conformation of the nascent chains, with their N termini facing the cytosol. This conformation is required for signal sequence processing. However, we discovered a class of Sec66-dependent proteins that engage only when they are long enough to adopt the looped conformation.

Finally, we monitored the fate of ER-associated ribosomes after translation termination using pulsed-labeling experiments. These data demonstrated that ER-associated ribosomes readily exchanged into the cytosol after at most a few rounds of translation at the ER.

CONCLUSION: These results, together with those in an accompanying Report on translation at mitochondria, establish proximity-specific ribosome profiling as a robust and general tool. In principle, this method can be applied to any site that can be specified by a biotin-ligase fusion protein. Thus, our approach provides *in vivo* access to a broad spectrum of subpopulations of ribosomes defined either by their subcellular locations or through their interactions with specific factors, such as chaperones. ■


RELATED ITEMS IN SCIENCE

C. C. Williams, C. H. Jan, J. S. Weissman, *Science* **346**, 1257522 (2014).
S. Shao, R. S. Hegde, *Science* **346**, 701–702 (2014).

Department of Cellular and Molecular Pharmacology, Howard Hughes Medical Institute, California Institute for Quantitative Biosciences, Center for RNA Systems Biology, University of California, San Francisco, San Francisco, CA 94158, USA.

*Corresponding author.

E-mail: jonathan.weissman@ucsf.edu
Cite this article as C. H. Jan, C. C. Williams, J. S. Weissman, *Science* **346**, 748–751 (2014).
DOI: 10.1126/science.1257521



Proximity-specific ribosome profiling provides spatiotemporal details of translation at the ER. Biotin ligase is localized to the ER as a fusion protein, where it biotinylates Avi-tagged ribosomes at the ER surface. Ribosome profiling is performed on streptavidin-purified ribosomes and compared to whole-cell profiling to resolve which genes are translated at the ER (1) and how much nascent chain was required to target the ribosome to the translocon (2).

RESEARCH ARTICLE

LOCAL TRANSLATION

Principles of ER cotranslational translocation revealed by proximity-specific ribosome profiling

Calvin H. Jan,* Christopher C. Williams,* Jonathan S. Weissman†

Localized protein synthesis is a fundamental mechanism for creating distinct subcellular environments. Here we developed a generalizable proximity-specific ribosome profiling strategy that enables global analysis of translation in defined subcellular locations. We applied this approach to the endoplasmic reticulum (ER) in yeast and mammals. We observed the large majority of secretory proteins to be cotranslationally translocated, including substrates capable of posttranslational insertion *in vitro*. Distinct translocon complexes engaged nascent chains at different points during synthesis. Whereas most proteins engaged the ER immediately after or even before signal sequence (SS) emergence, a class of Sec66-dependent proteins entered with a looped SS conformation. Finally, we observed rapid ribosome exchange into the cytosol after translation termination. These data provide insights into how distinct translocation mechanisms act in concert to promote efficient cotranslational recruitment.

Eukaryotic cells contain highly specialized subcellular environments, including both membrane- and nonmembrane-bound compartments. Localized protein synthesis can play a critical role in creating these subcellular structures by allowing protein production at the site of action and in response to local cellular need. Local translation is involved in diverse processes, including developmental patterning, cellular motility, synaptic plasticity, and protein trafficking through the secretory pathway (1). Dysfunctional RNA localization is linked to neurodevelopmental and neurodegenerative diseases (2). Numerous microscopy-based studies of individual mRNAs have demonstrated a breadth of subcellular localizations, and recent genome-wide mapping of transcript localization within cells and tissues has further emphasized the widespread spatial control of mRNA (3).

By contrast, global approaches for studying spatial control of protein synthesis are limited to bulk interrogations that cannot uniquely identify proteins—such as the RiboPuroMycylation (4) and FUNCAT (5) methods—or require careful biochemical fractionation of the compartment of interest (6), limiting both the location and resolution of analyses. These considerations motivated us to develop a generalizable strategy for enabling proximity-specific ribosome profiling that preserves *in vivo* spatiotemporal information about the site of synthesis. We employed a two-step

approach wherein we (i) used a spatially restricted biotin ligase (BirA) to mark ribosomes containing a biotin acceptor peptide (AviTag) in live cells with all membranes and spatial relations intact (7) and (ii) read out the translational activity of purified biotinylated ribosomes with ribosome profiling (the deep sequencing of ribosome-protected fragments) (8) that quantitatively reports on genome-wide translation with subcodon resolution (Fig. 1A).

Here we used this proximity-specific ribosome profiling strategy to study protein synthesis at the endoplasmic reticulum (ER), a major site of localized protein synthesis where a diverse set of proteins enter the secretory pathway. Work spanning several decades has revealed multiple routes of targeting nascent proteins to the ER (9). These include the canonical signal recognition particle (SRP)-dependent pathway in which translation is halted upon binding of SRP to hydrophobic sequences, and resumes only when the ribosome engages the translocon. Additionally, there are several SRP-independent pathways, although these are generally considered to mediate posttranslational import (9). Extensive studies have also elucidated the core translocational machinery necessary for protein import across and into the ER membrane, and identified accessory translocon factors in yeast and metazoans thought to increase the efficiency of protein import or assist the translocation of specific proteins (10).

Despite our in-depth mechanistic and structural understanding of these steps, the broader cellular organization of these targeting routes *in vivo* has remained largely unexplored. Experimental limitations have prevented a systematic characterization of substrate flux through

the various ER-targeting pathways in unperturbed cells. Similarly, our understanding of rough ER dynamics remains limited because of the difficulty in precisely measuring both the timing of ribosome-nascent chain (RNC) recruitment to the translocation machinery, as well as RNC fate following translation termination. Here we developed and applied proximity-specific ribosome profiling to address these fundamental questions.

A general approach for subcellular ribosome profiling: Development and application to the ER

To establish the proximity-specific ribosome profiling method, we implemented the following five steps: (i) introduction of a nonperturbing ribosome tag consisting of a tobacco etch virus (TEV) protease-cleavable AviTag; (ii) genetic targeting of BirA to a subcellular location of interest; (iii) temporal control of ribosome biotinylation *in vivo*; (iv) inhibition of postlysis biotinylation; and (v) selective isolation of biotinylated ribosomes and specific elution via TEV cleavage (Fig. 1A). We developed and validated these steps in the budding yeast *Saccharomyces cerevisiae*, as well as in the human embryonic kidney-293 (HEK-293) cell line.

Informed by a recent structure of the yeast 80S ribosome (11), we expressed Avi-tagged versions of several candidate ribosomal proteins with surface-accessible termini. We identified multiple subunits that when tagged and expressed from their endogenous loci, including the natural 3' untranslated region (UTR), were incorporated into ribosomes and covered growth defects seen in deletion mutants. These included C-terminally tagged *RPL16* and *RPS2* [also called uL13 and uS5 (12)], which were used for subsequent experiments (Fig. 1B). N-terminally tagged *RPL10a* [uL1] was used for mammalian studies (13) (fig. S1A).

For our yeast studies, we constructed three different ER-localized BirA fusion proteins, as well as cytosolic and mitochondrial controls (Fig. 1C). To broadly capture the translational activity of all ER-associated ribosomes, we localized BirA to the ER using the C-terminal tail-anchor (TA) from *UBC6* (14). To more specifically examine translation at two known translocation entry points to the ER, we fused BirA to *SEC63*, a member of the SEC complex that specifically associates with the Sec61 translocon (15), and to *SSH1*, a paralog of the canonical Sec61 translocon that interacts with SRP but not the SEC complex (16). For the mitochondrial studies, we used a BirA-fusion to *OM45*, a major constituent of the mitochondrial outer membrane (MOM). Finally, the mammalian studies used a BirA fusion to Sec61 β that uniformly labeled the ER (17). In all cases, the BirA fusion proteins showed the expected localization (Fig. 1C and fig. S1B).

Because of the potential cycling of ribosomes between different cellular locations, especially following translation termination, it was critical to be able to induce rapid ribosome biotinylation while also suppressing any constitutive background BirA activity. Although biotin is an essential

Department of Cellular and Molecular Pharmacology, Howard Hughes Medical Institute, California Institute for Quantitative Biosciences, Center for RNA Systems Biology, University of California, San Francisco, San Francisco, CA 94158, USA.

*These authors contributed equally to this work.

†Corresponding author. E-mail: jonathan.weissman@ucsf.edu

cofactor for both yeast and mammalian cells, titrating biotin levels in the growth media (7) suppressed BirA activity to undetectable levels without affecting cell growth (fig. S2). Brief biotin pulses were sufficient to give a robust biotinylation signal in live cells (Fig. 1E), and it was possible to prevent postlysis biotinylation by depleting lysates of biotin and adenosine 5'-triphosphate (ATP) (Fig. 1D and fig. S2). This procedure allowed us to achieve rapid (on the time scale of polypeptide synthesis) and efficient biotinylation of both our 40S and 60S Avi-tagged ribosomes using a cytosolic BirA. In marked contrast, ER-localized BirAs failed to label the 40S Avi-tagged ribosomal subunit but retained the ability to robustly label the 60S Avi-tagged subunit (Fig. 1E). Based on the length of our BirA tether, which is too short to allow biotinylation of the 40S subunit of a docked, translocating ribosome, this result demonstrates the specific biotinylation of oriented translocating ribosomes over those that passively encounter the ER membrane. Finally,

we optimized the purification of biotinylated, ribonuclease-digested monosomes (fig. S3) (18).

Validation of proximity-specific ribosome profiling

We performed proximity-specific ribosome profiling in *S. cerevisiae* using the three different ER-localized BirA constructs, as well as the cytosolic and mitochondrially localized controls (Fig. 2A), and in mammalian HEK-293 cells using an ER-localized BirA fusion protein. For each experiment, brief treatment with the translation elongation inhibitor cycloheximide (CHX), which preserves the ribosome position along an mRNA, was followed by a biotin pulse. Subsequent to processing and sequencing, we determined an enrichment value for each gene by taking the log₂ ratio of ribosome footprint densities in the matched streptavidin-pulldown versus input whole-cell ribosome profiling samples. Enrichment metrics obtained from the same BirA were highly reproducible between replicates (Fig. 2B; Ssh1 Pearson

$r = 0.97$; Sec63 Pearson $r = 0.98$) and robust across expression levels (fig. S4).

Targeting of BirA to the cytosol yielded a narrow range of enrichment values (90% of genes fell within -0.2 to $+0.2$ log₂ enrichment units), demonstrating that our protocol for isolating biotinylated monosomes introduced minimal bias. We detected a modest but significant ($P < 1 \times 10^{-15}$, Kolmogorov-Smirnov test) depletion of secreted genes, consistent with the expected lower accessibility of ER-docked ribosomes. By contrast, BirA targeted to mitochondria produced a clear bimodal distribution, enriching for genes annotated to localize to this cellular compartment (19). An in-depth analysis of translation at the mitochondrial outer membrane is presented in an accompanying Report (20). Targeting of BirA to the ER membrane inverted the mitochondrial enrichment pattern, cleanly separating secreted proteins from those synthesized in the cytosol or targeted to mitochondria. Ssh1, Sec63, and Ubc6 ER-localized BirA fusion


Fig. 1. A system for in vivo proximity-dependent ribosome biotinylation to monitor local protein synthesis at the ER.

(A) Schematic for proximity-specific ribosome profiling. (i) The *Escherichia coli* biotin ligase BirA is localized to a subcellular site of interest in cells expressing an Avi-tagged ribosomal protein and grown in low-biotin conditions. (ii) A biotin pulse is applied, resulting in specific biotinylation of ribosomes in close physical proximity to the localized BirA. (iii) Ribosome profiling of paired input (gray and red) and isolated biotinylated (red) monosomes reveals codon-resolved translational enrichment specific to the BirA locale.

(B) Fractionation of yeast lysates derived from strains containing scarless C-terminal Rps2 or Rpl16a/b hemagglutinin (HA)-TEV-AviTags on 10 to 50% sucrose gradients. Polysome traces demonstrate proper ribosomal assembly, and incorporation of tags into polysomes demonstrates their nonperturbative nature.

(C) ER localization of BirA fusion proteins used in this study. BirA-mVenus-Ubc6, Sec63-mVenus-BirA, and BirA-mVenus-Ssh1 all localize to the perinuclear and cortical ER. (D) Western blot analysis demonstrates that biotinylation of ribosomal AviTags does not occur before the addition of excess biotin or postlysis in our assay. (E)

Biotinylation kinetics of 40S and 60S AviTags by BirAs localized to the cytosol or ER (Sec63). Favorable kinetics were achieved independent of localization, and preferential 60S biotinylation demonstrates the specificity of the ER-localized ligase for oriented ER ribosomes. Shaded regions indicate biotinylation times used in subsequent sequencing experiments.



constructs labeled ribosomes translating similar sets of secretome genes [defined in (27)], though we observed pronounced differences in the point during translation at which RNCs interact with these BirA fusions (explored below). To determine whether other gene categories were significantly overrepresented in the enriched populations of our ER data sets, we performed gene ontology (GO-) term analysis on gene categories in yeast and mammalian cells that were enriched above a threshold derived from a receiver operator characteristic (ROC) analysis (Fig. 2, C and D, and fig. S5). In both yeast and HEK-293 cells, enriched gene sets were exclusively from the secretome (18). However, a substantial number of mammalian secretome transcripts predicted by Phobius (22) to encode secretory proteins were not enriched in our assay. This set of genes was enriched in GO-terms for nucleus and cytosol when compared to all secretome genes, arguing that these proteins represent potential false positives in the computationally predicted secretory gene set. This discrepancy serves to highlight both the sensitivity and utility of our approach for experimentally defining proteins that are targeted to specific cellular compartments.

We noted that peroxisomal proteins exhibited heterogeneous ER translational enrichment. The peroxisome is a highly conserved organelle responsible for lipid catabolism whose mechanism of biogenesis has been controversial. There is evidence for both de novo peroxisome generation from ER-derived vesicles, as well as for derivation from preexisting peroxisomes through growth and fission (23). Our data reveal that 16 of 54 yeast peroxisomal proteins showed clear cotranslational enrichment. Consistent with previous targeted studies in yeast (24), a unifying determinant for this ER targeting is the presence of one or more transmembrane domains (TMDs)—we found no evidence for the enrichment of peroxisomal matrix proteins (Fig. 2E). Notably, this partitioning was also seen in mammalian cells (fig. S6). Thus, peroxisomes appear to obtain transmembrane proteins from the ER and matrix proteins from the cytosol.

Cotranslational targeting of SRP-dependent and -independent substrates in vivo


Whereas a subset of proteins are strictly reliant on SRP for ER targeting, a process that is thought to be obligatorily cotranslational, import of other

proteins occurs efficiently without SRP when measured both in vitro and in vivo (21, 25). A recent in silico analysis revealed that ~40% of yeast secretome substrates use the less-studied SRP-independent pathway (27). SRP-independent translocation depends on translocon accessory factors, as well as the luminal chaperone Kar2/BiP, and in vitro can occur efficiently after translation (26, 27) (Fig. 3A).

Notably, we found that the vast majority of secretory proteins undergo cotranslational targeting in vivo, irrespective of their dependence on SRP (Fig. 3B). This pattern held across all BirA fusions (fig. S7) for 162 genes experimentally validated as SRP-dependent or -independent (21, 25), as well as for an additional 756 genes whose SRP dependence was predicted using a hydropathy-based analysis (27).

It was a formal possibility that the apparent cotranslational ER enrichment of these translating messages was a result of brief treatment with the translation elongation inhibitor (CHX) prior to biotinylation, because this provides extra time for the RNC complex to engage the translocon. We evaluated this possibility by omitting translation inhibitors and labeling with biotin

Fig. 2. Specificity of proximity-dependent ribosome profiling across multiple systems. (A) Boxplots of the log₂ enrichment distributions for secretome (blue), curated mitochondrial (red), and all other (gray) gene categories obtained from proximity-specific ribosome profiling experiments in yeast using different BirA fusions. Biotinylation was carried out in the presence of CHX for 2 min (cytosolic, mitochondria) or 7 min (ER). Enrichments were computed for each reliably expressed gene as the log₂ ratio of biotinylated footprint density (RPM) over the corresponding density from the matched input whole-cell ribosome profiling experiment.



for 1 min, a time scale comparable to that of a single round of polypeptide synthesis. For the large majority of SRP-dependent and -independent substrates, levels of translational enrichment were not dependent on CHX (Fig. 3B). Intriguingly, ribosomes translating a small minority of the SRP-independent proteins lost their enrichment, suggesting that in an unperturbed setting, these proteins translocate posttranslationally.

Thus, SRP independence is not synonymous with posttranslational translocation; import concurrent with protein synthesis is the principal route into the ER in vivo. By effectively coupling translation and translocation for the large majority of proteins entering the secretory pathway, the cell minimizes the dangers associated with having a cytosolic cohort of untranslocated, aggregation-prone proteins (28). Understanding how the cell achieves cotranslational translocation of SRP-independent messages remains unclear.

Comprehensive analysis of co- versus posttranslational translocation in vivo

Having uncoupled SRP independence from posttranslational translocation, we sought to better understand the determinants for partitioning between the co- and posttranslational import pathways. To classify genes on the basis of their ER translational enrichment, we systematically identified genes whose enrichments were dependent on CHX using a support vector machine (SVM) classifier trained to distinguish between ~140 proteins characterized empirically as being CHX-dependent or -independent (18). This SVM analysis enabled us to systematically characterize the import of proteins as being either cotranslational (CHX-independent), cotranslational translocation that is dependent upon (or enhanced by) treatment with a translation inhibitor (CHX-dependent), or obligatorily posttranslational (dis-enriched) (Fig. 3C). The enrichment of CHX-dependent proteins was greatest for Sec63-BirA, consistent with its role in co- and posttranslational translocation (fig. S8). These differences demonstrate the specificity of ribosome labeling by BirA fusion proteins and suggest that cotranslational insertion (CHX-independent) typically occurs through both translocons.

The SVM analysis indicated that the large majority (681 of 837) of Phobius-predicted secretory genes were translated at the ER independent of CHX. Of the remaining predicted secretory proteins, 63 were dependent on CHX for enrichment, whereas 93 were not enriched under any condition tested. The latter dis-enriched group contained nearly all of the roughly 50 annotated TA proteins whose C-terminal TMDs preclude cotranslational recognition and that are known to be targeted to the ER posttranslationally through the guided entry of TA proteins (GET) pathway (29, 30).

What, then, accounts for the remaining proteins whose translocation is not strictly cotranslational? The position of an ER-targeting signal within a protein imposes restrictions on when during synthesis targeting may occur, and thus might be an important determinant. Indeed,

robustly enriched CHX-independent secreted genes and dis-enriched TA genes fall on the opposite sides of this spectrum, with extreme N- and C-terminal ER-targeting signals, respectively (Fig. 3D). The targeting signals of proteins dependent on CHX for cotranslational targeting to the ER fall in between these two extremes, often present far downstream in relation to their over-

all gene length. Following cotranslational targeting to the ER, translation of long downstream regions leads to prolonged mRNA retention on the ER surface, which would be mediated by multiple translocating RNCs. In contrast, short downstream regions are unlikely to stably tether the mRNA to the ER following targeting (fig. S9). The CHX dependence of genes with short




Fig. 3. Global characterization of co- versus posttranslational translocation in vivo. (A) Overview of current models for SRP-dependent and -independent targeting to and translocation into the ER. Predictions for the proportion of substrates that partition between pathways are taken from (21). (B) Cumulative distribution of the Sec63-BirA \log_2 enrichments for SRP-dependent (blue), SRP-independent (red), and nonsecreted (gray) genes with or without CHX. Biochemically validated genes (dashed lines) were consolidated from (21) and (25). (C) Venn diagram summarizing the SVM classifications for CHX dependence in the context of the Phobius-predicted secretome. The Sec63-BirA +CHX enrichment profile was fit as a mixture of two normal distributions, and all genes enriched above the 99th percentile of the dis-enriched distribution were classified by the SVM. (D) Number of codons downstream of the first hydrophobic domain of Phobius-secretome genes versus the position of this domain relative to overall gene length, plotted for genes in different SVM-classified enrichment categories. Contour lines are added for specific gene sets for visual clarity and represent Gaussian density fits of the corresponding points in that set. Colors match those in (C) with the tail-anchored genes (dark-blue) overlaid as open circles. (E) Proportion of genes for which a hydrophobic feature was predicted by either TMHMM or SignalP, for different gene categories. Colors and gene sets match those in (C) with the addition of nonsecretome genes (gray), as predicted by Phobius.

downstream regions suggests that ER retention mediated by polysomes promotes efficient translocon targeting.

Our SVM classification also revealed cotranslational, CHX-independent ER enrichment of 70 genes for which no hydrophobic domains were detected by Phobius (Fig. 3C). However, most of these were predicted to contain a hydrophobic domain by alternate hydrophobic prediction algorithms (TMHMM or SignalP) (31, 32) (Fig. 3E). These genes thus likely represent genuine secretory proteins that were missed by Phobius, highlighting the value of our studies as an experimental complement to computational algorithms for globally identifying secreted and transmembrane proteins.

Timing and specificity of cotranslational targeting to the ER

We next asked when, during translation, RNCs are recruited to the ER, which is expected to de-

pend upon the mechanism of recruitment. For example, SRP binds preferentially to short nascent chains containing cytosolically accessible hydrophobic sequences (33) and halts translation elongation until the RNC reaches the ER. By contrast, SRP-independent transport through the SEC complex relies on a poorly understood network of cytosolic chaperones, none of which are known to arrest translation.

In yeast, translocation occurs through two paralogous channels, Sec61 and Ssh1. The essential Sec61 translocon associates with several accessory factors, including both essential (Sec63 and Sec62) and nonessential (Sec66 and Sec72) peripheral components to form the SEC complex, or separately with the SRP receptor (SR). By contrast, the nonessential Ssh1 is a simpler translocon thought to interact peripherally only with SR (16). We reasoned that fusing BirA to specific complexes would allow us to globally monitor the timing and specificity of translo-

cation of substrates through these distinct translocons (Fig. 4A).

Ssh1

Ssh1 is expected to receive RNCs from SRP and should therefore interact with the ribosomes after the hydrophobic sequence emerges from the ribosome peptide exit tunnel. Consistent with this model, for all secreted and TMD proteins, regardless of the location of their targeting sequence, BirA-Ssh1 enrichment began only after the hydrophobic domain was fully accessible (~60 amino acids from the start of targeting sequence) (34) (Fig. 4B and fig. S10A).

Type II signal anchors (SAs) and cleavable signal sequences (SSs) are oriented in a looped conformation with their N termini facing the cytosol (Fig. 4C). Models of how SSs and SAs achieve this topology within the translocon differ in the efficiency of RNC recruitment; i.e., the head-first model stipulates early RNC binding and subsequent




Fig. 4. Timing and specificity of co-translational targeting to the ER. (A) Schematic of the yeast translocon-specific BirAs used to examine ribosome accessibility at two translocational entry points into the ER. (B) Metagenote plots of \log_2 BirA-Ssh1 enrichment per codon (mean \pm SD) as a function of ribosome position relative to the first codon of the first Phobius-predicted hydrophobic element for the indicated signal class. Heat maps below represent single-gene positional

enrichments used to derive the corresponding averaged metagenote plot, sorted by increasing distance to the point at which enrichment occurs. (C) Violin plot showing the distribution of the point of enrichment for BirA-Ssh1 relative to the first hydrophobic element, for different types of hydrophobic features and Sec66-dependent genes as defined in (D). Shown above are two RNC conformations consistent with nascent chain lengths. (D) Gene enrichments obtained with the general BirA-Ubc6 ER marker in yeast in wild-type versus *sec66* Δ backgrounds. Sec66-dependent genes are defined in fig. S11. (E) Metagenote plot as in (B) of \log_2 BirA-Ubc6 enrichments for Sec66-dependent genes in wild-type (black) and *sec66* Δ (purple) backgrounds. (F) Metagenote plot as in (B) of \log_2 Sec63-BirA enrichments. (G) Metagenote plot of \log_2 enrichments as in (F) in a *sec55-1* SRP temperature-sensitive background at the permissive (25°C, black) and nonpermissive (37°C, red) temperatures.

signal inversion, whereas the looped-insertion model requires either delayed RNC binding in which the nascent chain is correctly oriented before binding (Fig. 4C) or delayed translocation following ribosome engagement. Studies of model SA substrates are consistent with the head-first model (35, 36). Efficient targeting to Ssh1 immediately after translation of the hydrophobic sequence is also consistent with such a mechanism. Here, the timing of engagement was bimodally distributed: Most of the secretome was efficiently recruited to Ssh1 immediately after exit of the hydrophobic sequence from the ribosome; however, a prominent subset of proteins with cleavable Ss engaged only after enough synthesis (~120 amino acids) to allow the nascent chain to acquire a looped topology (Fig. 4, B and C). Thus, both head-first and looped insertion can occur *in vivo*, depending on the protein. Deletion of *SEC66*, a nonessential component of the SEC complex that mediates translocation of SRP-independent substrates, exclusively affected translocation of the looped-insertion substrates (Fig. 4, C to E, and fig. S11). Thus, proximity-specific ribosome profiling can decipher how distinct translocon components enable the efficient handling of diverse targeting sequences and topologies (35, 37, 38).

Sec63

Sec63 mediates both SRP-independent and -dependent translocation. Without a strict require-

ment for SRP-induced translational pausing in the cytosol, Sec63 translocation of SRP-independent substrates naïvely might have been expected to result in a delayed and more broadly distributed timing of ER targeting. Surprisingly, the opposite was observed: Sec63 began to interact with RNCs translating secretory proteins well before the emergence of the hydrophobic sequence from the ribosome (Fig. 4F), and maximal engagement occurred shortly after this element was fully solvent accessible. Thus, Sec63 interacts with ribosomes through two distinct modes: One depends on the presence of an accessible hydrophobic element, and the second reflects an interaction with ribosomes while the targeting sequence is in the exit tunnel. Consistent with this interpretation, acute loss of SRP function using a temperature-sensitive SRP allele (fig. S12) did not affect the early engagement, but did compromise the late (exposed targeting sequence) enrichment (Fig. 4G).

Dynamics of ER-associated ribosomes

Upon translation termination, ER-associated ribosomes can either immediately dissociate and return to the pool of cytosolic ribosomes or preferentially undergo multiple rounds of translation on ER-associated mRNAs. To investigate these dynamics in the context of living cells, we harvested samples for proximity-specific profiling after increasing lengths of biotinylation time in the ab-

sence of CHX. Enrichment for secretory messages is expected to decrease at a rate proportional to the time scale at which biotinylated ribosomes, originating from the ER, exchange into the cytosol and begin translating cytosolic messages (Fig. 5A). We observed rapid collapse of our bimodal enrichment distribution into a single population on the order of minutes, although secreted messages remain on the enriched side of the distribution at all time points tested, as expected from continual biotinylation of ER ribosomes (Fig. 5B). Based on a median gene length of ~425 codons and a translation rate of ~5.5 codons per second (39), translation of a single secretory protein is expected to take ~77 s. Thus, ribosomes at the yeast ER are highly dynamic, freely exchanging into the cytosol within at most a few rounds of translation.

Discussion

Here we present a proximity-based ribosome profiling strategy that can monitor translation for any location at which it is possible to target a BirA fusion protein. We applied this strategy to analyze modes of cotranslational translocation into the ER. Nearly one-quarter of the proteome is imported into the ER; accordingly, this process has been the focus of intense research. Much of this previous work, however, has explored the behavior of a small group of model substrates, often outside of a cellular context. Proximity-specific ribosome profiling allowed us to simultaneously probe the ER engagement of nascent chains across the full proteome *in vivo*, in the context of competing and redundant targeting pathways. This comprehensive characterization revealed several principles of how cells integrate distinct targeting pathways with the translocation machinery to allow for unexpectedly robust cotranslational ER import of a diverse set of substrates.

Foremost is the critical role of the timing of translation of the ER-targeting sequence relative to translation termination for determining the propensity of a protein to undergo import cotranslationally. This stands in contrast to the view that cotranslational import is dictated by the factors that mediate targeting (e.g., SRP). It had previously been appreciated that TA proteins must insert posttranslationally because the targeting sequence is obscured prior to translation termination. However, these represent a single point on a broader spectrum of signal positions. Proteins with targeting domains near the C terminus typically were targeted as RNCs only when the kinetics of translation were crippled. By contrast, the predominantly SRP-independent (27) set of substrates containing N-terminal Ss are robustly cotranslationally targeted. Indeed, the full range of cotranslational substrates could engage both the essential SEC and “alternate” Ssh1 translocons. Moreover, RNCs were able to interact with the SEC complex prior to exposure of a hydrophobic sequence, even though there are no known mechanisms for coordinating translation and recruitment to SEC.

The above findings suggest a model wherein a pioneering round of translation is responsible




Fig. 5. Dynamics of ER-associated ribosomes in vivo. (A) Overview of the pulse-labeling experiment to assay the kinetics of ribosome exchange from the ER *in vivo*. (B) Histograms of log₂ Sec63-BirA enrichment values for well-expressed secretome (blue) and all other (gray) genes over the exchange time course. Times represent the total time of ribosome biotinylation in the absence of CHX. (C) Working model consistent with the positional enrichments observed for the translocon-specific BirAs and ribosome recycling. (1) Initial recruitment to the ER depends on a fully accessible signal sequence. (2) Ribosomes translating ER-tethered mRNAs can interact with SEC early. (3) Upon termination, ribosomes recycle into the cytosolic pool.

for recruiting the RNC to the ER surface, after which the message remains tethered to the ER by ongoing translation by downstream ribosomes (Fig. 5C). As evidenced by the lack of enrichment for proteins with low (i.e., length-limited) ribosome occupancy downstream of the hydrophobic targeting sequence, the tethered state appears to be crucial for efficient cotranslational engagement.

Consistent with our understanding of SRP function, SRP likely plays a critical role in establishing specificity and ensuring translocation competency through its ability to halt translation. Subsequent rounds of ribosome initiation in the context of this ER-tethered mRNA would obviate the need for SRP to survey every translation event, particularly for messages with extensive downstream regions that can accommodate multiple ribosomes. Such a mechanism is consistent with the observed 1:50 stoichiometry of SRP to the ribosome (40), and would simplify the problem of cellular protein sorting while minimizing the toxicity associated with solvent-exposed hydrophobic domains. Although initial recruitment may direct an RNC to a specific translocon, once tethered to the ER the high effective concentration would enable upstream RNCs to engage any translocon. Alternatively, the apparent lack of translocon substrate specificity could be due to ribosome biotinylation in trans, though the observed differences in position and CHX-dependent enrichments argue against this. In either case, inhibition of translocation is known to induce a massive cytoplasmic stress response (28), underscoring the danger of having ER-targeted proteins in the cytosol even when they can be posttranslationally translocated.

Our studies also revealed a class of SSs that emphasize an intimate connection between the timing of import and protein topology, mediated by translocon accessory factors. The bimodal timing of targeting to Ssh1 suggests that insertion can occur in either a head-first or looped orientation. Our results implicate Sec66 in mediating the import of those proteins that undergo looped insertion. This functionality may be necessary for certain substrates whose insertion kinetics would preclude reorientation within the translocon. A clear future application of our method is probing the elusive roles of other translocon accessory factors, such as the translocating chain-associated membrane protein (TRAM) and translocon-associated protein (TRAP) in mammals (37, 38).

A final principle that emerged from our studies is the dynamic nature of ER-associated ribosomes in yeast, which cycle readily between cellular compartments. This is in contrast to evidence from in vitro exchange experiments that showed stable association of the 60S ribosome subunit with the mammalian ER (41). It will be interesting to explore whether such a pattern holds in more specialized secretory cells, such as plasma cells, which rely on efficient translation at the ER. Indeed, electron micrographs have revealed the presence of circular polysomes (42) in these cells, consistent with a “closed-loop model” of trans-

lation (43) that is presumed to promote efficient translation reinitiation.

The principles uncovered here highlight the ability of proximity-specific ribosome profiling to synergize with prior mechanistic studies of ER-targeting pathways. Diverse biological systems localize mRNAs to generate cellular structure and function, yet compared to the ER, much less is known about how cotranslational protein targeting contributes to asymmetry at these sites. More generally, this approach enables the profiling of subpools of ribosomes that interact, even transiently, with cellular proteins of interest—e.g., those involved in protein folding, quality control, targeting, and posttranslational modification. As a flexible, precise, and global method, proximity-specific ribosome profiling provides a tool for exploring the interface between translation and cell biology.

REFERENCES AND NOTES

1. D. St Johnston, Moving messages: The intracellular localization of mRNAs. *Nat. Rev. Mol. Cell Biol.* **6**, 363–375 (2005). doi: [10.1038/nrm1643](#); pmid: [15852043](#)
2. H. Jung, C. G. Gkogkas, N. Sonenberg, C. E. Holt, Remote control of gene function by local translation. *Cell* **157**, 26–40 (2014). doi: [10.1016/j.cell.2014.03.005](#); pmid: [24679524](#)
3. J. H. Lee et al., Highly multiplexed subcellular RNA sequencing in situ. *Science* **343**, 1360–1363 (2014). doi: [10.1126/science.1250212](#); pmid: [24578530](#)
4. A. David et al., Nuclear translation visualized by ribosome-bound nascent chain purinification. *J. Cell Biol.* **197**, 45–57 (2012). doi: [10.1083/jcb.201112145](#); pmid: [22472439](#)
5. D. C. Dieterich et al., In situ visualization and dynamics of newly synthesized proteins in rat hippocampal neurons. *Nat. Neurosci.* **13**, 897–905 (2010). doi: [10.1038/nn.2580](#); pmid: [20543841](#)
6. S. B. Stephens, C. V. Nicchitta, In vitro and tissue culture methods for analysis of translation initiation on the endoplasmic reticulum. *Methods Enzymol.* **431**, 47–60 (2007). doi: [10.1016/S0076-6879\(07\)31004-5](#); pmid: [17932330](#)
7. M. Fernández-Suárez, T. S. Chen, A. Y. Ting, Protein-protein interaction detection in vitro and in cells by proximity biotinylation. *J. Am. Chem. Soc.* **130**, 9251–9253 (2008). doi: [10.1021/ja801445p](#); pmid: [18582056](#)
8. N. T. Ingolia, S. Ghaemmaghami, J. R. S. Newman, J. S. Weissman, Genome-wide analysis in vivo of translation with nucleotide resolution using ribosome profiling. *Science* **324**, 218–223 (2009). doi: [10.1126/science.1168978](#); pmid: [19213877](#)
9. T. Ast, M. Schuldiner, All roads lead to Rome (but some may be harder to travel): SRP-independent translocation into the endoplasmic reticulum. *Crit. Rev. Biochem. Mol. Biol.* **48**, 273–288 (2013). doi: [10.3109/10409238.2013.782999](#); pmid: [23530742](#)
10. E. C. Mandon, S. F. Trueman, R. Gilmore, Protein translocation across the rough endoplasmic reticulum. *Cold Spring Harb. Perspect. Biol.* **5**, a013342 (2013). doi: [10.1101/cshperspect.a013342](#); pmid: [23251026](#)
11. A. Ben-Shem et al., The structure of the eukaryotic ribosome at 3.0 Å resolution. *Science* **334**, 1524–1529 (2011). doi: [10.1126/science.1212642](#); pmid: [22096102](#)
12. N. Ban et al., A new system for naming ribosomal proteins. *Curr. Opin. Struct. Biol.* **24**, 165–169 (2014). doi: [10.1016/j.sbsi.2014.01.002](#); pmid: [24524803](#)
13. M. Heiman et al., A translational profiling approach for the molecular characterization of CNS cell types. *Cell* **135**, 738–748 (2008). doi: [10.1016/j.cell.2008.10.028](#); pmid: [19013281](#)
14. B. Kornmann et al., An ER-mitochondria tethering complex revealed by a synthetic biology screen. *Science* **325**, 477–481 (2009). doi: [10.1126/science.1175088](#); pmid: [19556461](#)
15. K. Finke et al., A second trimeric complex containing homologs of the Sec61p complex functions in protein transport across the ER membrane of *S. cerevisiae*. *EMBO J.* **15**, 1482–1494 (1996). pmid: [8612571](#)
16. Y. Jiang, Z. Cheng, E. C. Mandon, R. Gilmore, An interaction between the SRP receptor and the translocon is critical during cotranslational protein translocation. *J. Cell Biol.* **180**, 1149–1161 (2008). pmid: [18347066](#)
17. Y. Shibata et al., The reticulon and DPL/Yop1p proteins form immobile oligomers in the tubular endoplasmic reticulum. *J. Biol. Chem.* **283**, 18892–18904 (2008). doi: [10.1074/jbc.M800986200](#); pmid: [18442980](#)
18. Materials and methods are available as supplementary material on Science Online.
19. M. Elstner, C. Andreoli, T. Klopstock, T. Meitinger, H. Prokisch, The mitochondrial proteome database: MitoP2. *Methods Enzymol.* **457**, 3–20 (2009). doi: [10.1016/S0076-6879\(09\)05001-0](#); pmid: [19426859](#)
20. C. C. Williams, C. H. Jan, J. S. Weissman, Targeting and plasticity of mitochondrial proteins revealed by proximity-specific ribosome profiling. *Science* **346**, 748–751 (2014).
21. T. Ast, G. Cohen, M. Schuldiner, A network of cytosolic factors targets SRP-independent proteins to the endoplasmic reticulum. *Cell* **152**, 1134–1145 (2013). doi: [10.1016/j.cell.2013.02.003](#); pmid: [23452858](#)
22. L. Käll, A. Krogh, E. L. Sonnhammer, A combined transmembrane topology and signal peptide prediction method. *J. Mol. Biol.* **338**, 1027–1036 (2004). doi: [10.1016/j.jmb.2004.03.016](#); pmid: [15111065](#)
23. J. J. Smith, J. D. Aitchison, Peroxisomes take shape. *Nat. Rev. Mol. Cell Biol.* **14**, 803–817 (2013). doi: [10.1038/nrm3700](#); pmid: [24263361](#)
24. A. van der Zand, I. Braakman, H. F. Tabak, Peroxisomal membrane proteins insert into the endoplasmic reticulum. *Mol. Biol. Cell* **21**, 2057–2065 (2010). doi: [10.1091/mbc.E10-02-0082](#); pmid: [20427571](#)
25. D. T. Ng, J. D. Brown, P. Walter, Signal sequences specify the targeting route to the endoplasmic reticulum membrane. *J. Cell Biol.* **134**, 269–278 (1996). doi: [10.1083/jcb.134.2.269](#); pmid: [8707814](#)
26. B. P. Young, R. A. Craven, P. J. Reid, M. Willer, C. J. Stirling, Sec63p and Kar2p are required for the translocation of SRP-dependent precursors into the yeast endoplasmic reticulum in vivo. *EMBO J.* **20**, 262–271 (2001). doi: [10.1093/emboj/20.1.262](#); pmid: [11226176](#)
27. S. Panzner, L. Dreier, E. Hartmann, S. Kostka, T. A. Rapoport, Posttranslational protein transport in yeast reconstituted with a purified complex of Sec proteins and Kar2p. *Cell* **81**, 561–570 (1995). doi: [10.1016/0092-8674\(95\)90077-2](#); pmid: [7758110](#)
28. O. Brandman et al., A ribosome-bound quality control complex triggers degradation of nascent peptides and signals translation stress. *Cell* **151**, 1042–1054 (2012). doi: [10.1016/j.cell.2012.10.044](#); pmid: [23178123](#)
29. M. Schuldiner et al., The GET complex mediates insertion of tail-anchored proteins into the ER membrane. *Cell* **134**, 634–645 (2008). doi: [10.1016/j.cell.2008.06.025](#); pmid: [18724936](#)
30. S. Stefanovic, R. S. Hegde, Identification of a targeting factor for posttranslational membrane protein insertion into the ER. *Cell* **128**, 1147–1159 (2007). doi: [10.1016/j.cell.2007.01.036](#); pmid: [17382883](#)
31. A. Krogh, B. Larsson, G. von Heijne, E. L. Sonnhammer, Predicting transmembrane protein topology with a hidden Markov model: Application to complete genomes. *J. Mol. Biol.* **305**, 567–580 (2001). doi: [10.1006/jmbi.2000.4315](#); pmid: [11152613](#)
32. T. N. Petersen, S. Brunak, G. von Heijne, H. Nielsen, SignalP 4.0: Discriminating signal peptides from transmembrane regions. *Nat. Methods* **8**, 785–786 (2011). doi: [10.1038/nmeth.1701](#); pmid: [21959131](#)
33. V. Siegel, P. Walter, The affinity of signal recognition particle for presecretory proteins is dependent on nascent chain length. *EMBO J.* **7**, 1769–1775 (1988). pmid: [3169004](#)
34. M. Kowarik, S. Küng, B. Martoglio, A. Helenius, Protein folding during cotranslational translocation in the endoplasmic reticulum. *Mol. Cell* **10**, 769–778 (2002). doi: [10.1016/S1097-2765\(02\)00685-8](#); pmid: [12419221](#)
35. P. K. Devaraneni et al., Stepwise insertion and inversion of a type II signal anchor sequence in the ribosome-Sec61 translocon complex. *Cell* **146**, 134–147 (2011). doi: [10.1016/j.cell.2011.06.004](#); pmid: [21729785](#)
36. V. Goder, M. Spiess, Molecular mechanism of signal sequence orientation in the endoplasmic reticulum. *EMBO J.* **22**, 3645–3653 (2003). doi: [10.1093/emboj/cdg361](#); pmid: [12853479](#)
37. D. Görlich, E. Hartmann, S. Prehn, T. A. Rapoport, A protein of the endoplasmic reticulum involved early in polypeptide translocation. *Nature* **357**, 47–52 (1992). doi: [10.1038/357047a0](#); pmid: [1315422](#)
38. R. D. Fons, B. A. Bogert, R. S. Hegde, Substrate-specific function of the translocon-associated protein complex during translocation across the ER membrane. *J. Cell Biol.* **160**, 529–539 (2003). doi: [10.1083/jcb.200210095](#); pmid: [12578908](#)

39. B. Bonven, K. Gulløv, Peptide chain elongation rate and ribosomal activity in *Saccharomyces cerevisiae* as a function of the growth rate. *Mol. Gen. Genet.* **170**, 225–230 (1979). doi: [10.1007/BF00337800](https://doi.org/10.1007/BF00337800); pmid: [372763](https://pubmed.ncbi.nlm.nih.gov/372763/)
40. S. C. Ogg, P. Walter, SRP samples nascent chains for the presence of signal sequences by interacting with ribosomes at a discrete step during translation elongation. *Cell* **81**, 1075–1084 (1995). doi: [10.1016/S0092-8674\(05\)80012-1](https://doi.org/10.1016/S0092-8674(05)80012-1); pmid: [7600575](https://pubmed.ncbi.nlm.nih.gov/7600575/)
41. D. Borge, G. Blobel, D. D. Sabatini, In vitro exchange of ribosomal subunits between free and membrane-bound ribosomes. *J. Mol. Biol.* **74**, 415–438 (1973). doi: [10.1016/0022-2836\(73\)90037-5](https://doi.org/10.1016/0022-2836(73)90037-5); pmid: [4729519](https://pubmed.ncbi.nlm.nih.gov/4729519/)
42. A. K. Christensen, L. E. Kahn, C. M. Bourne, Circular polysomes predominate on the rough endoplasmic reticulum of somatotropes and mammotropes in the rat anterior pituitary. *Am. J. Anat.* **178**, 1–10 (1987). doi: [10.1002/aja.1001780102](https://doi.org/10.1002/aja.1001780102); pmid: [3825969](https://pubmed.ncbi.nlm.nih.gov/3825969/)
43. S. E. Wells, P. E. Hillner, R. D. Vale, A. B. Sachs, Circularization of mRNA by eukaryotic translation initiation factors. *Mol. Cell* **2**, 135–140 (1998). doi: [10.1016/S1097-2765\(00\)80122-7](https://doi.org/10.1016/S1097-2765(00)80122-7); pmid: [9702200](https://pubmed.ncbi.nlm.nih.gov/9702200/)

ACKNOWLEDGEMENTS

We thank members of the Weissman and P. Walter labs, as well as A. Ting, for discussion; and E. Costa, M. Kampmann, P. Kimmig, K. Kostova, and M. Smith for critical reading of the manuscript. Technical assistance was provided by the University of California, San Francisco, Center for Advanced Technology, Nikon Imaging Center, and the Consortium for Frontotemporal Dementia Research. This research was supported by the Center for RNA

Systems Biology and the Howard Hughes Medical Institute. C.H.J. is the Rebecca Ridley Kry Fellow of the Damon Runyon Cancer Research Foundation (DRG-2085-11). C.C.W. is an NSF Graduate Research Fellow. Data are deposited in Gene Expression Omnibus under accession nos. GSE61011 and GSE61012.

SUPPLEMENTARY MATERIALS

www.sciencemag.org/content/346/6210/1257521/suppl/DC1
Materials and Methods
Figs. S1 to S12
Tables S1 to S4
References (44–54)

16 June 2014; accepted 12 September 2014
10.1126/science.1257521